


Associations of serial negative income shock and all-cause mortality: a longitudinal study in China

Nan Xiao, Ran Li, Shanshan Li, Yudan Yu, Fan Yang, Jianan Yang, Gordon G Liu, Beini Lyu 

To cite: Xiao N, Li R, Li S, *et al*. Associations of serial negative income shock and all-cause mortality: a longitudinal study in China. *BMJ Public Health* 2024;**2**:e001512. doi:10.1136/bmjph-2024-001512

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/bmjph-2024-001512>).

Received 26 May 2024
Accepted 27 September 2024

ABSTRACT

Introduction With a precarious economic outlook and increasing income volatility in current times, understanding the association between negative income shock and health is crucial. However, few studies have examined such associations in developing countries. Using data from China, this study aimed to examine associations of both serial absolute income drops and relative income trajectory and mortality.

Methods We included 4757 participants from the China Health and Nutrition Survey, a large prospective cohort study. Data between 1989 and 1997 were used to define income drops and relative income trajectories. We defined income drop as a decrease of $\geq 50\%$ between two consecutive interviews and defined relative income trajectory using a latent class model. All-cause mortality between 2000 and 2015 was ascertained by participants' family members.

Results A total of 2066 (43.43%) experienced 1 income drop and 477 (10.03%) experienced ≥ 2 income drops. A total of 535 deaths occurred (incidence rate 8.88 per 1000 person-years). Income drops were associated with a greater risk of mortality after adjusting for baseline income, comorbidities, sociodemographic and behavioural factors (HR 1.42 (95% CI 1.04 to 1.93) for ≥ 2 income drops vs no income drop). The downshift in relative income was also associated with increased mortality risk (HR 3.61 (95% CI 1.45 to 8.96) for always low; HR 3.36 (95% CI 1.36 to 8.32) for decreasing; HR 2.92 (95% CI 1.14 to 7.51) for increasing vs always high relative income). The associations between income drops and mortality were observed only among individuals with low wealth and low household income.

Conclusion In a large sample of the Chinese population with repeated income measurement and over 14 years of follow-up, both serial absolute income drops and a downward relative income trajectory were associated with higher risks of mortality in China. Priority should be given to policies aimed at enhancing resilience against serial income shocks and financial burdens.

INTRODUCTION

Income volatility, specifically negative income shocks, has been associated with a higher risk of adverse outcomes such as depression,¹

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Although the association between negative income shock and health has been studied in various epidemiological literature, few studies have examined such associations in developing countries or examined long-term income changes.

WHAT THIS STUDY ADDS

⇒ This study demonstrates that independent of baseline income, both serial drops in absolute income and downshift in relative income were associated with higher risks of mortality. The associations were observed only among individuals with low wealth and low household income.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ These findings underscore the enduring significance of experiencing negative income shock as a predictor of mortality. In current times with increasing income volatility, these evidences call for strategies and policies to mitigate the health consequences of income drops.

cardiovascular disease² and all-cause mortality.³ Income volatility can occur due to factors such as seasonality of jobs, changes in work hours, voluntary or involuntary job changes and exits from work due to certain life events. Studies have found that more than one-third of US families experienced income volatility.^{4 5} Individuals and households with lower socioeconomic status are more likely to experience income volatility,^{6 7} which contributes to higher income inequality in the population.⁸

Although the association between negative income shock and health has been examined in many epidemiological studies, several research gaps remain. First, most prior literature originates from developed countries.^{3 9} The social welfare systems in developing countries, such as healthcare assistance and unemployment compensation, are



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. Published by BMJ.

Peking University, Beijing, China

Correspondence to
Dr Beini Lyu; blyu@pku.edu.cn

weaker, rendering individuals more vulnerable to negative income shock.¹⁰ In addition, developing countries generally have larger income inequality,¹¹ which is associated with worse health outcomes.^{12–13} Therefore, there is a need for more evidence from developing economies to inform efforts aimed at lowering health disparity. Second, the association between long-term income volatility and health outcomes was inadequately explored. Many studies measure income at a single time point, and only a few have examined income change between two time points.^{9–14–15} Incorporating multiple time points and providing a detailed characterisation of long-term income trajectories over the life course may better reflect individuals' time-varying financial status, which potentially has a stronger association with health.^{14–16–17} Third, most studies have solely focused on absolute income trajectory shift. However, relative deprivation in income rankings has been associated with worse health outcomes, independent of absolute income drops.^{18–20}

As a developing country, China is experiencing increasing income volatility.²¹ Using longitudinal data from the Chinese population with nearly two decades of follow-up, we aimed to examine the association between long-term income volatility and all-cause mortality in China, a country with rapid income changes and high-income inequality.²² While income volatility can arise from both decrease and increase in income, the current study focused on downward volatility because previous studies found that income drops had a greater impact on health.^{1–23} We examined the associations between both absolute income drops and changes in relative income and mortality.

METHODS

Study population

The China Health and Nutrition Survey (CHNS) is a prospective cohort study initiated in 1989. The CHNS sample was not designed to be nationally representative since CHNS has not obtained data to create a representative provincial-level sample by its inception.²⁴ Instead, the study sample was to be randomly selected to capture a range of economic and demographic circumstances. The original survey in 1989 used a multistage, random cluster process to draw the sample in each of eight provinces. Counties and cities in each province were stratified by income (low, middle and high) and four counties and two cities were selected in each province using a weighted sampling scheme. Villages and townships within the counties and urban and suburban neighbourhoods within the cities were also selected randomly. By 2011, the CHNS included 12 provinces and municipal cities, covering 288 communities, which accounted for 47% of China's population.²⁵ Participants have been followed up every 2–4 years with a total of 10 survey rounds by 2015.²⁶

The current study included participants who attended the 1989 or 1991 interview, participated in at least 3 interviews between 1989 and 1997 (a total of 4 surveys), were

aged 16–65 in 1989 and had complete income and history of comorbidity information between 1989 and 1997. Among 6340 participants, 4757 had complete mortality information by 2015 and were included in the final analysis, resulting in a lost to follow-up rate of 24.97% (see patient flow chart in online supplemental figure 1). Major causes of lost to follow-up included missing people (such as travel, hour of work or play), school children who were in boarding schools, migrant work for those aged above 16, and natural disaster and major redevelopment of housing in all large urban centres.²⁴

Additional information on the data and study materials is available on the CHNS official website (<https://www.cpc.unc.edu/projects/china>). This cohort study adhered to the Strengthening the Reporting of Observational Studies in Epidemiology reporting guideline.

Income predictor

Absolute income change

During CHNS interviews in the years 1989, 1991, 1993 and 1997, participants self-reported pretax individual income for the past 12 months, encompassing all sources, including business, farming, fishing, gardening, livestock, non-retirement wages and retirement pension. Reported incomes for each wave were inflated to RMB1989. An income drop was defined as a decrease of 50% or more in deflated income between two consecutive interviews. The predictor of interest in our analysis was the number of income drops between 1989 and 1997 (range 0–3), categorised into 0, 1 or ≥ 2 drops.

Relative income trajectory

We examined the trajectory of individuals' relative income over time. The relative income here denotes individuals' relative position or rank compared with others. We grouped individuals based on their income quartile at each interview between 1989 and 1997 to measure their relative income hierarchy within the population. A latent class model was employed to identify subgroups characterised by similar underlying trajectories across income quartiles over the study period. This model used the maximum likelihood method to estimate the posterior probabilities of membership for each participant in each trajectory class. Participants were assigned to the class with the highest probability of membership.^{27–29}

Mortality

Our primary outcome was all-cause mortality. Participants who were alive in 1997 were followed from the year 2000 until death, lost to follow-up or the last interview in 2015, whichever occurred first. In the CHNS, mortality status was updated in each survey round. Participants' family members provided information on whether they were alive or deceased, and if deceased, the year of death.

Covariates

Covariates were selected if previous studies suggested potential associations with both income shock and mortality.^{2–3} Baseline characteristics collected in 1989 or

1991 included age at enrolment, ethnicity, sex, completion of middle school education, occupation (farmer, civil servant, self-employment or others), health insurance status and *hukou* status. In China, all households have to be registered in the locale where they reside and their *hukou* status is categorised as either 'rural' or 'urban'. Residents with urban *hukou* enjoy better health insurance, housing, pensions and other privileges, and rural-to-urban migration was heavily restricted. Additionally, we included participants' self-reported smoking status (yes/no), alcohol consumption (yes/no) and physical activity. Physical activity (yes/no) was determined by whether participants reported spending time weekly in six categories of physical activity (martial arts, running, gymnastics, soccer or basketball, badminton or volleyball and ping pong). In addition, we included the history of comorbidities (ie, myocardial infarction, stroke, diabetes and hypertension) and overweight. Overweight was defined as a body mass index greater than 24 kg/m² in the Asian context,³⁰ calculated as weight in kilograms divided by height in metres squared.

Categorical covariates that could vary over time, such as health conditions, were defined at baseline between 1989 and 1997. For comorbidities, smoking, alcohol consumption and overweight status, participants were categorised based on whether they ever exhibited the condition or behaviour during the period from 1989 to 1997.

Statistical analysis

We presented participants' baseline characteristics between 1989 and 1997 by the number of income drops and by relative income trajectory classes. Continuous variables were presented as mean (SD) or median (IQR) and categorical variables were presented as number (percentage) as appropriate. We used global χ^2 tests to compare categorical variables and analysis of variance to compare continuous variables across income categories.

We compared baseline characteristics between participants lost to follow-up and those included in analyses to examine potential impact of lost to follow-up on results. For the latent class model in the relative income analysis, the model was fitted with 2–6 classes, and the best fitting model was selected based on criteria including Akaike information criterion (AIC), Bayesian information criterion (BIC) and entropy. We calculated the incidence rate of all-cause mortality (per 1000 person-years) by income predictor category. Log-rank test was used to compare unadjusted Kaplan-Meier curves across categories of income predictors. We then assessed the association between income change and all-cause mortality using Cox proportional hazards models. The proportional hazard assumption was tested by checking the Schoenfeld's partial residuals and no violation was found. We fit an unadjusted model, a model adjusted for sociodemographic characteristics and logarithm of 1997 income, and a model further adjusted for lifestyle factors and self-reported history of comorbidities. All data analyses were

conducted with Stata, V.17 MP. A two-sided $p < 0.05$ was considered statistically significant.

Subgroup analyses

Given the potential buffering effect of wealth and household income on individual income shocks, we stratified individuals by the median of baseline wealth and household income. Additionally, we explored heterogeneity in the associations between income shock and mortality by age and sex.

Secondary analyses

We conducted the following sensitivity analyses to validate our results. First, we re-examined the association between income drops and mortality using alternative 25% and 75% drops in absolute income as cut points. Second, because income may change over time, we adjusted for time-dependent income levels as covariates from 2000 to 2015 to account for the impact of income on mortality. Third, because individuals living in one household may share resources and household income volatility could also be associated with risk of mortality, we replicated our analyses using household per capita income.

To explore potential mechanisms, we further conducted several secondary analyses. First, we assessed whether the effect of income shock on mortality was mediated by health-related factors by comparing behaviour profiles among participants with different numbers of income drops. We constructed a healthy lifestyle score which included smoking, alcohol consumption and physical activity. Each indicator was assigned 1 point for a healthy behaviour (never smoking, no alcohol consumption and ever being physically active) and 0 points for unhealthy alternatives. The healthy lifestyle score was the sum of the points and ranged from 0 to 3, with higher scores indicating healthier lifestyles. Second, we assessed whether the effect of income shock on mortality was mediated by healthcare access. As there is no direct measurement of healthcare accessibility in CHNS, we defined individuals reporting higher illness frequency but lower medical expenditure as having limited healthcare access. Illness frequency was defined as the number of reported sickness or injuries in the last 4 weeks during the follow-up period, while medical expenditure encompassed total treatment costs incurred when seeking medical care. Associations between illness frequency, medical expenditure and income drops were assessed separately using an analysis of variance.

RESULTS

Baseline characteristics

Among the 4757 participants, the mean (SD) age was 37.68 (12.67) years, with 2512 (52.81%) being men. Of these participants, 2214 (46.54%) experienced no income drop at baseline, 2066 (43.43%) experienced 1 income drop and 477 (10.03%) experienced ≥ 2 income drops (table 1). Compared with participants with no income drops, individuals with income drops were more likely

Table 1 Baseline characteristics of participants by number of income drops, CHNS

	Income drops, n				P value
	Overall	0	1	≥2	
Number of individuals	4757	2214	2066	477	
Age, years	37.68 (12.67)	38.29 (12.15)	37.32 (13.15)	36.38 (12.78)	<0.001
Male, n (%)	2512 (52.81)	1152 (52.03)	1084 (52.47)	276 (57.86)	0.06
Ethnicity, Han, n (%)	4156 (87.37)	1928 (87.08)	1811 (87.66)	417 (87.42)	0.85
<i>Hukou</i> , urban, n (%)	1120 (23.54)	686 (30.98)	363 (17.57)	71 (14.88)	<0.001
Education, completion of middle school, n (%)	1629 (34.24)	817 (36.90)	670 (32.43)	142 (29.77)	<0.001
Health insurance, n (%)	1793 (37.69)	1038 (46.88)	641 (31.03)	114 (23.90)	<0.001
Occupation, n (%)					
Farmer	3154 (66.30)	1224 (55.28)	1536 (74.35)	394 (82.60)	<0.001
Civil servant	1339 (28.15)	882 (39.84)	394 (19.07)	63 (13.21)	<0.001
Self-employment	158 (3.32)	71 (3.21)	75 (3.63)	12 (2.52)	0.43
Other	106 (2.23)	37 (1.67)	61 (2.95)	8 (1.68)	0.01
1997 income, RMB, median (IQR)	2819 (1233–4919)	2765 (1385–4527)	2780 (1064–5142)	3538 (1034–6978)	<0.001
Ever smoker, n (%)	2167 (45.58)	975 (44.04)	958 (46.37)	234 (49.06)	0.09
Ever drinker, n (%)	932 (19.60)	444 (20.05)	389 (18.84)	99 (20.75)	0.48
Ever exercise, n (%)	400 (8.41)	205 (9.26)	168 (8.13)	2 (5.66)	0.03
Ever overweight, n (%)	1020 (21.45)	512 (23.13)	406 (19.66)	102 (21.38)	0.02
Comorbidities, n (%)					
Myocardial infarction	113 (2.38)	62 (2.80)	45 (2.18)	6 (1.26)	0.10
Stroke	164 (3.45)	82 (3.70)	67 (3.24)	15 (3.14)	0.66
Diabetes	216 (4.54)	124 (5.60)	76 (3.68)	16 (3.35)	<0.001
Hypertension	257 (5.40)	128 (5.78)	117 (5.67)	12 (2.52)	0.01

Continuous variables were presented as mean (SD). Categorical variables were presented as number (percentage). CHNS, China Health and Nutrition Survey.

to be younger, less likely to have urban *hukou*, complete middle school or have health insurance ($p < 0.001$ for all). Individuals experiencing income drops were more likely to be farmers and less likely to be civil servants. Furthermore, they had a lower prevalence of overweight, diabetes and hypertension. No significant difference was observed in other characteristics.

Participants included in the current study were similar to those lost to follow-up for most demographic characteristics (online supplemental table 1). However, participants lost to follow-up were more likely to have urban *hukou*, be civil servants and less likely to be farmers.

Associations between income drops and all-cause mortality

A total of 535 death events occurred between 2000 and 2015, with a median follow-up time of 14 (IQR 9–18) years, resulting in 60 232 person-years. Participants with ≥ 2 income drops were at significantly greater risk of mortality compared with those with no income drop (log-rank $p < 0.05$, figure 1A). The incidence rate for all-cause mortality increased with the number of income drops (10.43 (95% CI 8.14 to 13.36) per 1000 person-years for individuals with ≥ 2 income drops vs 8.47 (95% CI 7.46 to 9.62) per 1000 person-years for those

with 0 drop, table 2). In the Cox proportional model, compared with participants with no income drops, those with ≥ 2 income drops had greater unadjusted hazards of all-cause mortality, although the association was not statistically significant (HR 1.23; 95% CI 0.93 to 1.63). After adjusting for baseline income, demographic, socio-economic status, lifestyle and history of comorbidities, participants with ≥ 2 drops had a significantly higher risk of mortality (HR 1.42 (95% CI 1.04 to 1.93)). We did not find a significant difference in risk of mortality between participants with one income drop and those with 0 drop (HR 1.06 (95% CI 0.86 to 1.32)).

Associations between relative income trajectory and all-cause mortality

In the latent class analysis of relative income change between 1989 and 1997, we found that the model best fit with four classes based on the values of AIC, BIC, adjusted BIC, entropy and p value of the Vuong-Lo-Mendell-Rubin likelihood test (online supplemental table 2 and figure 2). The four groups were categorised as follows: always high income (271 participants, 5.70%), increasing income (730, 15.35%), decreasing income (1042, 21.90%) and always low income (2714, 57.05%).




Figure 1 Kaplan-Meier curves for all-cause mortality by the number of income drops (A) and by relative income trajectory groups (B).

Compared with individuals with always high relative income, those with decreasing and always low relative income were older, and less likely to have urban *hukou* ($p<0.001$ for all, online supplemental table 3). They were also less likely to have health insurance, to be civil

servants, to drink alcohol, to smoke and to be overweight ($p<0.001$ for all).

Participants with always low income had the highest mortality risk, followed by those with decreasing, increasing and always high income (figure 1B). Compared with always high income, always low income, decreasing income and increasing income were associated with significantly higher risks of mortality (unadjusted HR=3.67 (95% CI 1.90 to 7.12) for always low; HR=3.12 (95% CI 1.58 to 6.16) for decreasing and HR=2.29 (95% CI 1.13 to 4.64) for increasing, table 3). After adjusting for baseline income and other potential confounders, the associations remained (compared with always high, adjusted HR=3.61 (95% CI 1.45 to 8.96) for always low; adjusted HR=3.36 (95% CI 1.36 to 8.32) for decreasing; adjusted HR=2.92 (95% CI 1.14 to 7.51) for increasing). Individuals with increasing income were at lower risks of mortality when compared with those with decreasing relative income (adjusted HR=0.84 (95% CI 0.54 to 1.31)) and always low relative income (adjusted HR=0.82 (95% CI 0.55 to 1.23)), but the associations were not statistically significant.

Subgroup analyses

In subgroup analyses, the association between ≥ 2 income drops and mortality was only significant among those who had lower wealth (HR 1.90; 95% CI 1.21 to 2.99 for lower wealth vs HR 1.04; 95% CI 0.68 to 1.61 for higher wealth, p for interaction=0.02, figure 2). Similarly, the association was only significant among those with lower household income (HR 1.56; 95% CI 1.04 to 2.34 for lower household income vs HR 1.30; 95% CI 0.79 to 2.12 for higher household income, p for interaction=0.02). In contrast, the associations between ≥ 2 income drops and mortality were similar among male and female participants. Furthermore, the association between ≥ 2 income drops was slightly lower among those who aged over 50 (HR 1.34; 95% CI 0.87 to 2.09) compared with their younger counterparts (HR 1.55; 95% CI 1.00 to 2.39), but

Table 2 Incidence and risk of all-cause mortality by income groups

	n (%)	No. of events/ person-years	IR* (95% CI)	Model 1 HR (95% CI)	Model 2 HR (95% CI)	Model 3 HR (95% CI)
Overall	4757	535/60 232	8.88 (8.16 to 9.66)	–	–	–
Income drops, n (%)						
0	2214 (46.54)	236/27 857	8.47 (7.46 to 9.62)	1 (Ref)	1 (Ref)	1 (Ref)
1	2066 (43.43)	237/26 430	8.97 (7.90 to 10.18)	1.07 (0.89 to 1.28)	1.09 (0.88 to 1.34)	1.06 (0.86 to 1.32)
≥ 2	477 (10.03)	62/5945	10.43 (8.14 to 13.36)	1.23 (0.93 to 1.63)	1.43 [†] (1.05 to 1.94)	1.42 [†] (1.04 to 1.93)

Model 1 is unadjusted. Model 2 is adjusted for 1989 age, gender, ethnicity, *hukou* status, completion of middle school education, health insurance status, occupation and logarithm of 1997 income. Model 3 is additionally adjusted for smoking status, alcohol consumption, physical activity, overweight status, reported history of myocardial infarction, stroke, diabetes and hypertension.

*Per 1000 person-years.

[†]Statistically significant, $p<0.05$.

IR, incidence rate.

Table 3 Incidence and risk of all-cause mortality by income trajectory classes

	n (%)	No. of events/ person-years	IR* (95% CI)	Model 1 HR (95% CI)	Model 2 HR (95% CI)	Model 3 HR (95% CI)
Overall	4757	535/60 232	8.88 (8.16 to 9.66)	–	–	–
Income trajectory classes, n (%)						
Always high	271 (5.70)	9/3337	2.70 (1.40 to 5.18)	1 (Ref)	1 (Ref)	1 (Ref)
Increasing	730 (15.35)	53/8628	6.14 (4.70 to 8.03)	2.29 [†] (1.13 to 4.64)	2.97 [†] (1.16 to 7.61)	2.92 [†] (1.14 to 7.51)
Decreasing	1042 (21.90)	110/12 941	8.50 (7.06 to 10.24)	3.12 [†] (1.58 to 6.16)	3.32 [†] (1.34 to 8.19)	3.36 [†] (1.36 to 8.32)
Always low	2714 (57.05)	363/35 326	10.28 (9.28 to 11.38)	3.67 [†] (1.90 to 7.12)	3.58 [†] (1.45 to 8.87)	3.61 [†] (1.45 to 8.96)

Model 1 is unadjusted. Model 2 is adjusted for 1989 age, gender, ethnicity, hukou status, completion of middle school education, health insurance status, occupation and logarithm of 1997 income. Model 3 is additionally adjusted for smoking status, alcohol consumption, physical activity, overweight status, reported history of myocardial infarction, stroke, diabetes and hypertension.
 *Per 1000 person-years.
[†]Statistically significant, p<0.05.
 IR, incidence rate.

the difference was statistically insignificant (p for interaction=0.15).

Sensitivity analyses

When using 25% and 75% as cut-offs for income drops, the patterns of mortality and different numbers of income drops remained consistent (online supplemental table 4). In fully adjusted models, the association between ≥2 income drops and mortality was stronger when using 75% as cut-off (HR 1.47; 95% CI 0.90 to 2.41) compared with 50% cut-off and 25% cut-off (HR 1.37; 95% CI 1.00 to 1.88).

After adjusting for time-dependent income between 2000 and 2015, the association between income drops

and mortality remained materially unchanged (online supplemental table 5). Similar to the main results, both serial absolute household per capita income drops and a low relative household per capita income trajectory were associated with significantly higher risks of mortality (online supplemental figure 3, online supplemental tables 6 and 7).

Compared with participants with no income drops, those with ≥2 income drops had lower healthy lifestyle scores during the follow-up period (online supplemental table 8, mean 1.56; 95% CI 1.49 to 1.63 for ≥2 drops vs 1.66; 95% CI 1.62 to 1.69 for 0 drops). Although we found no differences in the probability of illness during




Figure 2 Subgroup analysis of income drops and all-cause mortality. Model adjusted for 1989 age, gender, ethnicity, hukou status, completion of middle school education, health insurance status, occupation, logarithm of 1997 income, smoking status, alcohol consumption, physical activity, overweight status, reported history of myocardial infarction, stroke, diabetes and hypertension.

the follow-up period among different income drop groups (0.16; 95% CI 0.11 to 0.21 for ≥ 2 drops vs 0.18; 95% CI 0.16 to 0.20 for 0 drops, $p > 0.1$), individuals with ≥ 2 income drops had significantly lower medical expenditure compared with those with no income drops (RMB436; 95% CI RMB26 to RMB846 for ≥ 2 drops vs RMB793; 95% CI RMB467 to RMB1118 for 0 drops, $p = 0.03$).

DISCUSSION

In a large sample of the Chinese population with repeated income measurement and over 14 years of follow-up, we found that independent of baseline income, both serial absolute income drops and a low relative income trajectory during early to mid-adulthood were associated with a higher risk of mortality. Notably, these associations were observed only among individuals with lower wealth and household income, and potentially more impactful among younger people. These findings underscore the enduring significance of experiencing negative income shock as a predictor of mortality.

As a developing country, China has experienced rapid growth in income with massive structural shifts in economy. In the 1980–1990s, China introduced economic reforms and started to transform from a planned economy to market-oriented economy. Together with other significant policy changes, such as reforms to the pension system and the hardening of budget constraints on state enterprises, these changes have increased uncertainty and income volatility.²¹ In our study, we found that farmers and people with lower socioeconomic status were more likely to experience income drops, which is consistent with previous studies.^{6 31} The higher-income volatility among farmers could be potentially attributed to several reasons. For example, farmers may have unstable product outlets due to a market-oriented economy. Additionally, their income was vulnerable to natural disaster such as a series of severe floods along the Yangtze River and Pearl River in China during the 1990s.^{32 33}

The associations found in our study were consistent with previous studies conducted in high-income countries.^{2 3 34} For instance, a study using Norwegian register data found that males aged 50–69 who experienced at least two substantial falls in income had a 17% excess mortality rate.³⁴ Similarly, an analysis based on a cohort of young adults in the USA showed that experiencing more than two income drops over a 15-year period was associated with a 1.92-fold increase in the risk of death.³ Another study using a US sample from the Panel Study of Income Dynamics found that income level was a strong predictor of mortality, especially for persons under the age of 65 years.³⁵ In the current study, we found that ≥ 2 income drops over 14 years were associated with 1.42-fold increased risk of all-cause mortality. Sensitivity analyses showed that income drop using 75% cut-off had stronger association with mortality when compared with analyses using 25% or 50% cut-off. However, the association was

not statistically significant when using 75% cut-off, likely due to the smaller sample size and a smaller number of events among those with two income drops using 75% cut-off.

Until now, most studies examining the association between income changes and all-cause mortality have been conducted in high-income, developed countries. Despite the weaker social welfare systems and increased vulnerability of populations in developing countries, there has been a notable absence of research addressing the association between serial income shocks and mortality within these contexts. A study using the China Kadoorie Biobank found that individuals experiencing sharp socioeconomic status downshift between baseline and resurvey had a significant 30% reduction in survival time before cardiovascular deaths.³⁶ However, like other studies on income and health outcomes,^{10 37–39} it was unable to capture serial income changes across multiple time points. Our study examined longer-term income volatility and showed that independent of baseline income, serial drops in absolute income and low relative income were associated with higher risks of mortality. In addition, we found a dose–response association between income drops and mortality, with more income drops associated with greater mortality risk.

Our study revealed that the association between negative income shock and all-cause mortality was observed only among individuals with low-wealth and low-household income. Specifically, among participants with high baseline wealth and household income, ≥ 2 income drops were not associated with risks of mortality compared with no income drop. The results suggested that higher wealth and household income may mitigate some impacts of income drops on these mechanisms and weaken the association with mortality. In sensitivity analyses, we found that lifestyle and healthcare accessibility are potential mechanisms underlying the association between income drop and mortality. This is consistent with previous studies suggesting that income drops may induce stress-coping behaviours such as increased cigarette smoking⁴⁰ and alcohol consumption,⁴¹ longer working hours at the expense of physical activities,^{42 43} as well as delayed medical care and incomplete adherence to prescription medications.⁴⁴

Interestingly, we found that increasing relative income was associated with a slightly lower risk of mortality compared with individuals with always low income and decreasing income, though the association was not statistically significant. This may suggest that upward relative income change could to some extent modify the greater risk of mortality among individuals with low baseline income. An upshift in socioeconomic status has been associated with a lower risk of cardiovascular mortality in a Korean study.⁴⁵ Potential mechanisms could be that rising socioeconomic status results in healthier lifestyle,⁴⁶ better affordability of healthcare services⁴⁷ and better support from members in the same socioeconomic class.⁴⁸

With a precarious economic outlook and increasing income volatility in current times, our study underscores the importance of prioritising policies aimed at enhancing resilience against serial income shocks. In the USA, unemployment and wage insurance have been implemented as short-term strategies to offset the burdens of transitory income shocks such as job loss.⁴⁹ Unemployment insurance might reduce the adverse health consequences of job loss by shielding workers from financial stress,⁵⁰ improving access to care⁵¹ and also diminishing the psychosocial impact of unemployment.⁵² Studies have shown that countries with more generous unemployment insurance systems exhibit better health outcomes and narrower health inequalities.^{53–55} In China, some local governments have launched vocational skills subsidy programmes to equip recently unemployed individuals with the necessary skills to find jobs.⁵⁶ By addressing the root causes of income instability and providing targeted support to vulnerable populations, policy-makers can help mitigate the detrimental effects of income volatility on health and well-being.

Our study had several strengths. First, previous studies were predominantly conducted in developed countries, while our study focused on a large developing country. We confirmed that in developing countries with weaker social welfare systems, independent of baseline income, long-term drops in income were associated with increased mortality risk. Moreover, we highlighted the independent impact of relative income changes on survival risk. Second, the repeated income data in the CHNS allowed us to capture detailed income change and the long follow-up time allowed us to assess the mortality risk over nearly two decades. Third, rather than relying solely on theoretical frameworks, we explored potential mechanisms underlying negative income shock and mortality using population data. Our findings underscore the pivotal roles of health behaviours and healthcare access in mediating these associations.

However, we acknowledge several limitations in our study. First, there was no standard definition of substantial income change. Prior studies have used 25% and 75% as the cut-off for drop.^{3 57} Nevertheless, the associations in our study remained consistent across various thresholds (50%, 25% and 75%). Similar to Elfassy *et al*,³ our analysis noted that more than two 25% drops in income were substantial enough to significantly impact mortality risk. Second, studies on income change and health are subject to reverse causation, where individuals who become ill may be more prone to job loss and subsequent negative income shock. We controlled for baseline comorbidities and were able to preserve temporal order by using income change information between 1989 and 1997 to assess mortality in the subsequent follow-up. Third, self-reported income data might be susceptible to recall bias. Fourth, participants included in our study were likely more socioeconomically advantaged compared with those lost to follow-up, which may bias our estimates of the associations.

Conclusions

Using data from a developing country, we found that independent of baseline income, both serial absolute income drops and a low relative income trajectory during early to mid-adulthood were associated with a higher risk of mortality. We also found that wealth and household income could mitigate some of the adverse effects. These findings underscore the enduring significance of experiencing negative income shock as a predictor of mortality. In current times with increasing economic uncertainty and income volatility, these evidences call for policies to mitigate the health consequences of income shocks.

Acknowledgements This research uses data from China Health and Nutrition Survey (CHNS). We are grateful to research grant funding from the National Institute for Health (NIH), the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD) for R01 HD30880 and R01 HD38700, National Institute on Aging (NIA) for R01 AG065357, National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) for R01 DK104371 and P30 DK056350, National Heart, Lung and Blood Institute (NHLBI) for R01 HL108427, the NIH Fogarty grant D43 TW009077, the Carolina Population Center for P2C HD050924 and P30 AG066615 since 1989 and the China-Japan Friendship Hospital, Ministry of Health for support for CHNS 2009, Chinese National Human Genome Center in Shanghai since 2009 and Beijing Municipal Center for Disease Prevention and Control since 2011. We thank the National Institute for Nutrition and Health, China Center for Disease Control and Prevention, Beijing Municipal Center for Disease Control and Prevention and the Chinese National Human Genome Center in Shanghai.

Contributors NX and BL conceptualised the study. NX conducted the literature search, performed the data collection and analysis. NX and BL had directly accessed and verified the underlying data. BL supervised the project administration. NX and BL completed the first draft of the manuscript. NX, RL, SL, YY, FY, JY, GGL and BL interpreted the results and critically revised the manuscript. NX, RL, SL, YY, FY, JY, GGL and BL read and approved the submitted version, had full access to all the data in the study and accepted responsibility to submit for publication. BL is the guarantor.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and the CHNS was approved by the institutional review boards at the University of North Carolina at Chapel Hill and the National Institute of Nutrition and Health of the Chinese Center for Disease Control and Prevention (ethical approval number: 2015-024). Each CHNS participant has provided a written informed consent. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. This article uses data from the China Health and Nutrition Survey (CHNS), which are available to the scientific community on submitting a registration form to the CHNS study.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is

properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Beini Lyu <http://orcid.org/0000-0001-9219-468X>

REFERENCES

- Prause J, Dooley D, Huh J. Income Volatility and Psychological Depression. *Am J Comm Psychol* 2009;43:57–70.
- Wang SY, Tan ASL, Claggett B, et al. Longitudinal Associations Between Income Changes and Incident Cardiovascular Disease: The Atherosclerosis Risk in Communities Study. *JAMA Cardiol* 2019;4:1203–12.
- Elfassy T, Swift SL, Glymour MM, et al. Associations of Income Volatility With Incident Cardiovascular Disease and All-Cause Mortality in a US Cohort. *Circulation* 2019;139:850–9.
- Hannagan A, Morduch J. Income Gains and Month-to-Month Income Volatility: Household Evidence from the Us Financial Diaries. *SSRN Journal* 2015.
- Burhouse S, Chu K, Ernst K, et al. FDIC National Survey of Unbanked and Underbanked Households. 2016. Available: <https://www.fdic.gov/household-survey/2015-fdic-national-survey-unbanked-and-underbanked-households>
- Yang Y, Zhang S. Understanding income volatility in urban China. *J Asian Econ* 2023;85:101588.
- Hardy BL. Childhood Income Volatility and Adult Outcomes. *Demography* 2014;51:1641–65.
- Nichols A. Income inequality, volatility, and mobility risk in China and the US. *China Econ Rev* 2010;21:S3–11.
- Boen C. The role of socioeconomic factors in Black-White health inequities across the life course: Point-in-time measures, long-term exposures, and differential health returns. *Soc Sci Med* 2016;170:63–76.
- Luo W, Xie Y. Socio-economic disparities in mortality among the elderly in China. *Popul Stud (Camb)* 2014;68:305–20.
- Chancel L, Piketty T, Saez E, et al. *World inequality report 2022*. Harvard University Press.
- Auger N, Zang G, Daniel M. Community-level income inequality and mortality in Québec, Canada. *Public Health (Fairfax)* 2009;123:438–43.
- Park J, Ryu S-Y, Han M, et al. The association between income inequality and all-cause mortality across urban communities in Korea. *BMC Public Health* 2015;15:574.
- Thomson RM, Igelström E, Purba AK, et al. How do income changes impact on mental health and wellbeing for working-age adults? A systematic review and meta-analysis. *Lancet Public Health* 2022;7:e515–28.
- Frech A, Damaske S. Men's Income Trajectories and Physical and Mental Health at Midlife. *Am J Sociol* 2019;124:1372–412.
- Beydoun MA, Beydoun HA, Mode N, et al. Racial disparities in adult all-cause and cause-specific mortality among us adults: mediating and moderating factors. *BMC Public Health* 2016;16:1113.
- de Mestral C, Stringhini S. Socioeconomic Status and Cardiovascular Disease: an Update. *Curr Cardiol Rep* 2017;19:115.
- Åberg Yngwe M, Fritzell J, Lundberg O, et al. Exploring relative deprivation: Is social comparison a mechanism in the relation between income and health? *Soc Sci Med* 2003;57:1463–73.
- Subramanyam M, Kawachi I, Berkman L, et al. Relative deprivation in income and self-rated health in the United States. *Soc Sci Med* 2009;69:327–34.
- Kondo N, Saito M, Hikichi H, et al. Relative deprivation in income and mortality by leading causes among older Japanese men and women: AGES cohort study. *J Epidemiol Community Health* 2015;69:680–5.
- Chamon M, Liu K, Prasad E. Income uncertainty and household savings in China. *J Dev Econ* 2013;105:164–77.
- Xie Y, Zhou X. Income inequality in today's China. *Proc Natl Acad Sci USA* 2014;111:6928–33.
- Grasset L, Glymour MM, Elfassy T, et al. Relation between 20-year income volatility and brain health in midlife: The CARDIA study. *Neurology (Echronicon)* 2019;93:e1890–9.
- Popkin BM, Du S, Zhai F, et al. Cohort Profile: The China Health and Nutrition Survey—monitoring and understanding socio-economic and health change in China, 1989–2011. *Int J Epidemiol* 2010;39:1435–40.
- Zhang B, Zhai FY, Du SF, et al. The China Health and Nutrition Survey, 1989–2011. *Obes Rev* 2014;15:2–7.
- Zhang Z, Liu M, Zhou C, et al. Evaluation of Dietary Niacin and New-Onset Hypertension Among Chinese Adults. *JAMA Netw Open* 2021;4:e2031669.
- Kendzor DE, Caughy MO, Owen MT. Family income trajectory during childhood is associated with adiposity in adolescence: a latent class growth analysis. *BMC Public Health* 2012;12:611.
- Grant RW, McCloskey J, Hatfield M, et al. Use of Latent Class Analysis and k-Means Clustering to Identify Complex Patient Profiles. *JAMA Netw Open* 2020;3:e2029068.
- Wickwire EM, Albrecht JS, Capaldi VF 2nd, et al. Trajectories of Insomnia in Adults After Traumatic Brain Injury. *JAMA Netw Open* 2022;5:e2145310.
- Tian Y, Jiang C, Wang M, et al. BMI, leisure-time physical activity, and physical fitness in adults in China: results from a series of national surveys, 2000–14. *Lancet Diabetes Endocrinol* 2016;4:487–97.
- Whalley J, Yue X. Rural Income Volatility and Inequality in China. *CESifo Econ Stud* 2009;55:648–68.
- Dao-yi G, Jin-hong Z, Shao-wu W. Flooding 1990s along the Yangtze River, has it concern of global warming? *J Geogr Sci* 2001;11:43–52.
- Zhi-Yong W, Gui-Hua L, Zhi-Yu L, et al. Trends of Extreme Flood Events in the Pearl River Basin during 1951–2010. *Adv Clim Change Res* 2013;4:110–6.
- Bævre K, Kravdal Ø. The effects of earlier income variation on mortality: An analysis of Norwegian register data. *Popul Stud (Camb)* 2014;68:81–94.
- McDonough P, Duncan GJ, Williams D, et al. Income dynamics and adult mortality in the United States, 1972 through 1989. *Am J Public Health* 1997;87:1476–83.
- Lai R, Li R, Wang T, et al. Short-term socioeconomic status shift and long-term cardiovascular outcomes in China: a population-based cohort study. *J Epidemiol Community Health* 2023;77:152–9.
- Li H, Zhu Y. Income, income inequality, and health: Evidence from China. *J Comp Econ* 2006;34:668–93.
- Yu X, Zhang W. All-cause mortality rate in China: do residents in economically developed regions have better health? *Int J Equity Health* 2020;19:12.
- Zhao Z. Income Inequality, Unequal Health Care Access, and Mortality in China. *Pop & Dev Rev* 2006;32:461–83.
- Kendzor DE, Businelle MS, Costello TJ, et al. Financial Strain and Smoking Cessation Among Racially/Ethnically Diverse Smokers. *Am J Public Health* 2010;100:702–6.
- Shaw BA, Agahi N, Krause N. Are changes in financial strain associated with changes in alcohol use and smoking among older adults? *J Stud Alcohol Drugs* 2011;72:917–25.
- Kochar A. Smoothing Consumption by Smoothing Income: Hours-of-Work Responses to Idiosyncratic Agricultural Shocks in Rural India. *Rev Econ Stat* 1999;81:50–61.
- Paffenbarger RS Jr, Hyde RT, Wing AL, et al. Physical activity, all-cause mortality, and longevity of college alumni. *N Engl J Med* 1986;314:605–13.
- Heisler M, Choi H, Rosen AB, et al. Hospitalizations and deaths among adults with cardiovascular disease who underuse medications because of cost: a longitudinal analysis. *Med Care* 2010;48:87–94.
- Sung J, Song YM, Hong KP. Relationship between the shift of socioeconomic status and cardiovascular mortality. *Eur J Prev Cardiol* 2020;27:749–57.
- Zhang Y-B, Chen C, Pan X-F, et al. Associations of healthy lifestyle and socioeconomic status with mortality and incident cardiovascular disease: two prospective cohort studies. *BMJ* 2021;373:n604.
- Rahimi AR, Spertus JA, Reid KJ, et al. Financial barriers to health care and outcomes after acute myocardial infarction. *JAMA* 2007;297:1063–72.
- Chandola T. Social inequalities in health by individual and household measures of social position in a cohort of healthy people. *J Epidemiol Community Health* 2003;57:56–62.
- Mitman K, Rabinovich S. Optimal unemployment insurance in an equilibrium business-cycle model. *J Monet Econ* 2015;71:99–118.
- Renahy E, Mitchell C, Molnar A, et al. Connections between unemployment insurance, poverty and health: a systematic review. *Eur J Public Health* 2018;28:269–75.
- Berkowitz SA, Basu S. Unemployment Insurance, Health-Related Social Needs, Health Care Access, and Mental Health During the COVID-19 Pandemic. *JAMA Intern Med* 2021;181:699–702.
- Young C. Losing a Job: The Nonpecuniary Cost of Unemployment in the United States. *Soc Forces* 2012;91:609–34.
- Ferrarini T, Nelson K, Sjöberg O. Unemployment insurance and deteriorating self-rated health in 23 European countries. *J Epidemiol Community Health* 2014;68:657–62.

- 54 Norström T, Grönqvist H. The Great Recession, unemployment and suicide. *J Epidemiol Community Health* 2015;69:110–6.
- 55 Cylus J, Glymour MM, Avendano M. Do generous unemployment benefit programs reduce suicide rates? A state fixed-effect analysis covering 1968–2008. *Am J Epidemiol* 2014;180:45–52.
- 56 Liang Y, Yu L. Human capital and the re-employment of retrenchment labor in urban China. *Rev Dev Econ* 2019;23:1432–58.
- 57 Pool LR, Burgard SA, Needham BL, *et al.* Association of a Negative Wealth Shock With All-Cause Mortality in Middle-aged and Older Adults in the United States. *JAMA* 2018;319:1341–50.